

# Week 14 – Partial Fractions Improper Integrals

---

## Partial Fractions Expansion

Given an algebraic expression like

$$\frac{3}{x-2} + \frac{5}{x+4}$$

we can write it with a common denominator as:

$$\frac{3}{x-2} + \frac{5}{x+4} = \frac{8x+2}{(x-2)(x+4)}$$

To evaluate the integrals like

$$\int \frac{(8x+2)}{(x-2)(x+4)} dx$$

we have to reverse this process.

$$\frac{8x+2}{(x-2)(x+4)} = \frac{A}{x-2} + \frac{B}{x+4}$$

$$8x+2 = A(x+4) + B(x-2)$$

$$x=2 \Rightarrow A = \frac{18}{6} = 3$$

$$x=-4 \Rightarrow B = \frac{-30}{-6} = 5$$

Now, the integral can easily be evaluated in terms of logarithms.

For a given rational function  $R(x) = \frac{P(x)}{Q(x)}$ , keep in mind the following:

- If degree of  $P(x)$  is greater than the degree of  $Q(x)$ , divide them using polynomial division.
- If  $Q(x)$  contains a power of the type  $(ax+b)^n$ , include all the terms

$$\frac{A_1}{ax+b} + \frac{A_2}{(ax+b)^2} + \dots + \frac{A_n}{(ax+b)^n}$$

in the expansion.

- An irreducible second order polynomial  $x^2+bx+c$  can be transformed into  $u^2 \pm a^2$  with the substitution  $u = x + \frac{b}{2}$ .

**Exercise 14-1:** Evaluate  $\int \frac{4x+4}{(x-3)(x-2)} dx$

**Solution:**

$$\frac{4x+4}{(x-3)(x-2)} = \frac{A}{x-3} + \frac{B}{x-2}$$

$$4x+4 = A(x-2) + B(x-3)$$

$$x=3 \Rightarrow A = 16$$

$$x=2 \Rightarrow B = -12$$

$$\int \left( \frac{16}{x-3} - \frac{12}{x-2} \right) dx$$

$$= 16 \ln |x - 3| - 12 \ln |x - 2| + c$$

**Exercise 14-2:** Evaluate

$$\int \frac{5x^3 - 12x^2 - 6x - 5}{x^2 - 2x - 3} dx$$

**Solution:** First, we have to make a polynomial division to obtain:

$$\frac{5x^3 - 12x^2 - 6x - 5}{x^2 - 2x - 3} = 5x - 2 + \frac{5x - 11}{(x - 3)(x + 1)}$$

$$\frac{5x - 11}{(x - 3)(x + 1)} = \frac{A}{x - 3} + \frac{B}{x + 1}$$

$$5x - 11 = A(x + 1) + B(x - 3)$$

$$x = 3 \Rightarrow A = 1$$

$$x = -1 \Rightarrow B = 4$$

$$\begin{aligned} \int \left( 5x - 2 + \frac{1}{x - 3} + \frac{4}{x + 1} \right) dx \\ = \frac{5}{2}x^2 - 2x + \ln |x - 3| + 4 \ln |x + 1| + c \end{aligned}$$

**Exercise 14-3:** Evaluate

$$\int \frac{10x^2 - 22x + 7}{x(x - 1)^2} dx$$

**Solution:**

$$\frac{10x^2 - 22x + 7}{x(x - 1)^2} = \frac{A}{x} + \frac{B}{x - 1} + \frac{C}{(x - 1)^2}$$

$$10x^2 - 22x + 7 = A(x - 1)^2 + Bx(x - 1) + Cx$$

Solving these equations, we obtain:

$$A = 7, \quad B = 3, \quad C = -5$$

$$\begin{aligned} \int \left( \frac{7}{x} + \frac{3}{x - 1} - \frac{5}{(x - 1)^2} \right) dx \\ = 7 \ln |x| + 3 \ln |x - 1| + \frac{5}{x - 1} + c \end{aligned}$$

**Exercise 14-4:** Evaluate

$$\int \frac{8x^2 + 11x + 18}{x^3 + 4x^2 + x + 4} dx$$

**Solution:**

$$\begin{aligned} \frac{8x^2 + 11x + 18}{x^3 + 4x^2 + x + 4} &= \frac{8x^2 + 11x + 18}{(x^2 + 1)(x + 4)} \\ &= \frac{Ax + B}{x^2 + 1} + \frac{C}{x + 4} \end{aligned}$$

$$8x^2 + 11x + 18 = Ax(x + 4) + B(x^2 + 1) + C(x + 4)$$

Solving these equations, we obtain:

$$A = 2, \quad B = 3, \quad C = 6$$

$$\begin{aligned} \int \left( \frac{2x + 3}{x^2 + 1} + \frac{6}{x + 4} \right) dx \\ = \ln |x^2 + 1| + 3 \arctan x + 6 \ln |x + 4| + c \end{aligned}$$

**Exercise 14-5:** Evaluate  $\int \frac{dx}{x^2 + 4x + 5}$

**Solution:** The substitution  $u = x + 2$  gives:

$$x^2 + 4x + 5 = u^2 + 1, \quad du = dx$$

$$\begin{aligned} \int \frac{1}{x^2 + 4x + 5} dx &= \int \frac{1}{u^2 + 1} du \\ &= \arctan u + c \\ &= \arctan(x + 2) + c \end{aligned}$$

**Exercise 14-6:** Evaluate the following integrals:

a)  $\int \frac{dx}{x^2 + x - 6}$

b)  $\int \frac{6x^3 - 18x}{(x^2 - 1)(x^2 - 4)} dx$

c)  $\int \frac{4x^4 + x + 1}{x^5 + x^4} dx$

d)  $\int \frac{1}{4x^2 + 4x - 3} dx$

e)  $\int \frac{3x + 1}{(x^2 + 2x + 5)^2} dx$

f)  $\int \frac{1}{\sqrt{3 - 2x - x^2}} dx$

## Improper Integrals

The integral

$$\int_a^b f(x) dx$$

exists if  $[a, b]$  is a closed and bounded interval and  $f$  is a continuous function. If the interval is unbounded like

$$[a, \infty), (-\infty, +\infty)$$

or if the function  $f$  has an infinite discontinuity, then the integral is called an improper integral. We define improper integrals by a limit process.

**Improper Integrals of type I:** Assuming the function  $f$  is continuous on the region of integration:

- $\int_a^{\infty} f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$
- $\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$
- $\int_{-\infty}^{+\infty} f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^{+\infty} f(x) dx$

If those limits exist, we say the integral converges, otherwise, we say the integral diverges.

**Exercise 14-7:** Evaluate  $\int_1^{\infty} \frac{dx}{x^2}$

**Solution:**

$$\begin{aligned} \int_1^{\infty} \frac{dx}{x^2} &= \lim_{t \rightarrow \infty} \int_1^t \frac{dx}{x^2} \\ &= \lim_{t \rightarrow \infty} -\frac{1}{x} \Big|_1^t \\ &= \lim_{t \rightarrow \infty} 1 - \frac{1}{t} \\ &= 1 \end{aligned}$$

So the integral is convergent.

**Exercise 14-8:** Evaluate  $\int_1^{\infty} \frac{dx}{\sqrt{x}}$

**Solution:**

$$\begin{aligned} \int_1^{\infty} \frac{dx}{\sqrt{x}} &= \lim_{t \rightarrow \infty} \int_1^t \frac{dx}{\sqrt{x}} \\ &= \lim_{t \rightarrow \infty} 2\sqrt{x} \Big|_1^t \\ &= \lim_{t \rightarrow \infty} 2\sqrt{t} - 2 \\ &= \infty \end{aligned}$$

So the integral is divergent.

**Improper Integrals of type II:**

- If  $f$  is continuous on  $(a, b]$  and discontinuous at  $a$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$$

- If  $f$  is continuous on  $[a, b)$  and discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

- If  $f$  is continuous on  $[a, b]$  except for  $c$  where  $a < c < b$  then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

**Example:** Evaluate  $\int_0^2 \frac{dx}{(x-1)^2}$

**Solution:**

$$\int_0^2 \frac{dx}{(x-1)^2} = \int_0^1 \frac{dx}{(x-1)^2} + \int_1^2 \frac{dx}{(x-1)^2}$$

$$= \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{(x-1)^2} + \lim_{t \rightarrow 1^+} \int_t^2 \frac{dx}{(x-1)^2}$$

$$\lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{(x-1)^2} = \lim_{t \rightarrow 1^-} -\frac{1}{x-1} \Big|_0^t$$

$$= \lim_{t \rightarrow 1^-} -\frac{1}{t-1} - 1$$

$$= \infty$$

So the integral is divergent. Note that

$$\int_0^2 \frac{dx}{(x-1)^2} = -\frac{1}{x-1} \Big|_0^2$$

gives an incorrect result.

**Direct Comparison Test:** Suppose that  $f$  and  $g$  are continuous on  $[a, \infty)$  and

$$0 \leq f(x) \leq g(x)$$

for all  $x \in [a, \infty)$ .

- If  $\int_a^\infty g(x) dx$  converges, then  $\int_a^\infty f(x) dx$  also converges.
- If  $\int_a^\infty f(x) dx$  diverges, then  $\int_a^\infty g(x) dx$  also diverges.

**Exercise 14-9:** Is the integral  $\int_1^\infty \frac{dx}{x^2 + x^5}$  convergent?

**Solution:** It is difficult to evaluate this integral, but we don't have to. We know that

$$x > 1 \Rightarrow x^2 + x^5 > x^2$$

$$\Rightarrow \frac{1}{x^2 + x^5} < \frac{1}{x^2}$$

$\int_1^\infty \frac{dx}{x^2}$  is convergent, therefore  $\int_1^\infty \frac{dx}{x^2 + x^5}$  is also convergent by comparison test.

**Limit Comparison Test:** Suppose that positive functions  $f$  and  $g$  are continuous on  $[a, \infty)$  and

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L$$

where  $0 < L < \infty$ , then the integrals

$$\int_a^\infty f(x) dx \quad \text{and} \quad \int_a^\infty g(x) dx$$

both converge or both diverge.

**Exercise 14-10:** Is the integral  $\int_1^\infty \frac{5+2x}{3+x^2} dx$  convergent?

**Solution:** It is difficult to evaluate this integral, but we don't have to. We know that

$$\lim_{x \rightarrow \infty} \frac{\frac{5+2x}{3+x^2}}{\frac{1}{x}} = 2$$

$\int_1^\infty \frac{dx}{x}$  is divergent, therefore  $\int_1^\infty \frac{5+2x}{3+x^2} dx$  is also divergent by limit comparison test.

## Review Exercises

**Exercise 14-11:** Evaluate the following integrals:

a)  $\int \frac{x^2}{x^4 - 1} dx$

b)  $\int \frac{x^4}{x^2 + 4} dx$

c)  $\int \frac{dx}{x^3 + x}$

d)  $\int \frac{dx}{\sqrt{5 + 12x - 9x^2}}$

e)  $\int x\sqrt{3 + 2x - x^2} dx$

**Exercise 14-12:** Evaluate the following integrals if they converge:

a)  $\int_1^\infty \frac{1}{x^2} dx$

b)  $\int_0^1 \frac{1}{\sqrt{x}} dx$

c)  $\int_1^\infty \frac{1}{x^p} dx$

d)  $\int_1^4 \frac{dx}{(x-2)^{2/3}}$

e)  $\int_3^4 \frac{dx}{(x-3)^2}$

f)  $\int_{-\infty}^{+\infty} x^3 dx$

**Exercise 14-13:** Determine the convergence or divergence of the following integrals:

a)  $\int_0^\infty \frac{1}{x + e^x} dx$

b)  $\int_0^\infty e^{-x^2} dx$

c)  $\int_1^\infty \frac{2 + \sin x}{\sqrt{x}} dx$

d)  $\int_1^{\infty} \frac{x^2 e^x}{\ln x} dx$

e)  $\int_{\pi}^{\infty} \frac{dx}{\sqrt{x} - \sin x}$

## Answers

### Exercise 14-11:

a)  $\frac{1}{2} \arctan x + \frac{1}{4} \ln |x - 1| - \frac{1}{4} \ln |x + 1| + c$

b)  $\frac{x^3}{3} - 4x + 8 \arctan \frac{x}{2} + c$

c)  $\ln |x| - \frac{1}{2} \ln |x^2 + 1| + c$

d)  $\frac{1}{3} \arcsin \left( x - \frac{2}{3} \right) + c$

e)  $\frac{1}{6} (2x^2 - x - 9) \sqrt{3 + 2x - x^2} + 2 \arcsin \frac{x - 1}{2} + c$

### Exercise 14-12:

a) 1

b) 2

c)  $\frac{1}{p - 1}$  if  $p > 1$

d)  $3 + 3\sqrt[3]{2}$

e) Divergent

f) Divergent

### Exercise 14-13:

a) Convergent

b) Convergent

c) Divergent

d) Divergent

e) Divergent

— End of WEEK —

**Author:** Dr. Emre Sermutlu

**Last Update:** December 22, 2016