

Week 6 – Derivative

The slope of the curve $y = f(x)$ at the point x_0 is:

$$m = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

if the limit exists. The tangent line through the point $(x_0, f(x_0))$ has this slope.

Derivative: The derivative of the function $f(x)$ is the function $f'(x)$ defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

We can think of the derivative as the rate of change of a function f or the slope of the curve of $y = f(x)$.

We will use

$$y', \quad f'(x), \quad \frac{dy}{dx}, \quad \frac{d}{dx}f(x)$$

to denote derivatives and

$$f'(a), \quad \left. \frac{dy}{dx} \right|_{x=a}$$

to denote their values at a certain point. Note that derivative is a function, its value at a point is a number.

Example: Instantaneous velocity is defined as

$$v = \lim_{\Delta t \rightarrow 0} \frac{x(t + \Delta t) - x(t)}{\Delta t}$$

In mechanics, velocity is the derivative of distance and acceleration is the derivative of velocity.

Exercise 6-1: Find the derivative of $f(x) = x^2$ using the definition.

Solution: $f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h}$

$$\begin{aligned} &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} \\ &= \lim_{h \rightarrow 0} 2x + h \\ &= 2x \end{aligned}$$

Exercise 6-2: Find the derivative of $f(x) = \sqrt{x}$ using the definition.

Solution: $f'(x) = \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h}$

$$\begin{aligned} &= \lim_{h \rightarrow 0} \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})} \\ &= \lim_{h \rightarrow 0} \frac{1}{\sqrt{x+h} + \sqrt{x}} \\ &= \frac{1}{2\sqrt{x}} \end{aligned}$$

Exercise 6-3: Find the derivative of $f(x) = \frac{1}{x}$ using the definition.

Differentiation Formulas: Using the definition of derivative, we obtain:

• Derivative of a constant is zero, i.e. $\frac{dc}{dx} = 0$

• Derivative of $f(x) = x$ is

$$f'(x) = 1$$

• Derivative of $f(x) = x^2$ is

$$f'(x) = 2x$$

• Derivative of $f(x) = x^n$ (where n is a positive integer) is:

$$f'(x) = nx^{n-1}$$

• Derivative of $f(x) = \sqrt{x}$ is:

$$f'(x) = \frac{1}{2\sqrt{x}}$$

- If f is a function and c is a constant, then

$$(cf)' = cf'$$

- If f and g are functions, then

$$(f + g)' = f' + g'$$

(Derivative is a linear operator)

Exercise 6-4: Evaluate the derivatives of the following functions:

a) $f(x) = 7x^3 + 3x^2 - 8$

b) $f(x) = 2\sqrt{x} - \frac{1}{4}x^4$

c) $f(x) = \frac{7x^3 - 18x + \sqrt{x}}{x}$

Exercise 6-5: Find the equation of the line that is tangent to $y = x^8 + x^4 + x^2$ at the point $(1, 3)$

Exercise 6-6: Find a horizontal tangent to

$$y = x^3 + x^2 - x + 1$$

Exercise 6-7: Find the equation of the tangent line to the graph at the given point. Then, sketch the function and the tangent line on the same coordinate system:

a) $f(x) = \frac{1}{x}$ at $(1, 1)$

b) $f(x) = \sqrt{x}$ at $(9, 3)$

c) $f(x) = x^3$ at $(0, 0)$

d) $f(x) = x^2$ at $(1, 1)$

e) $f(x) = x^3$ at $(1, 1)$

Differentiability and Continuity: If f is differentiable at a , then it is continuous at a , but if it is continuous at a , it is not necessarily

differentiable.

If the graph of a function has no unique tangent at a point, then the function has no derivative at that point. For example, $f(x) = |x|$ is not differentiable at $x = 0$. (Its derivative does not exist)

Higher Order Derivatives: We can find the derivative of the derivative of a function. It is called second derivative and denoted by:

$$y'', \quad f''(x), \quad \frac{d^2y}{dx^2}$$

For third derivative, we use f''' but for fourth and higher derivatives, we use the notation $f^{(4)}(x)$

Exercise 6-8: Let $f(x) = 7x^3 - 18x$. Find f' , f'' and f''' .

Vertical Tangents: The curve $y = f(x)$ has a vertical tangent at $(a, f(a))$ if f is continuous at a and if $|f'(x)| \rightarrow \infty$ as $x \rightarrow a$.

Exercise 6-9: Find the equation of the tangent line to $f(x) = \sqrt{x}$ at $(0, 0)$ and sketch the function and the tangent line on the same coordinate system.

Product Rule: If f and g are differentiable at x , then fg is differentiable at x and

$$\frac{d}{dx}(fg) = \frac{df}{dx}g + f\frac{dg}{dx}$$

or more briefly:

$$(fg)' = f'g + fg'$$

Reciprocal Rule: If f is differentiable at x and if $f(x) \neq 0$ then:

$$\left(\frac{1}{f}\right)' = \frac{-f'}{f^2}$$

Quotient Rule: If f and g are differentiable at x , and $g(x) \neq 0$ then $\frac{f}{g}$ is differentiable at x and:

$$\left(\frac{f}{g}\right)' = \frac{f'g - g'f}{g^2}$$

Exercise 6-10: Evaluate the derivatives of the following functions:

a) $f(x) = \frac{x^2 - 2}{x^2 + 2}$

b) $f(x) = \frac{x^4 - 7x^2 + 5}{x}$

c) $f(x) = \frac{1}{1 - x^{-3}}$

d) $f(x) = \frac{x + \sqrt{x}}{x^2 - \sqrt{x}}$

Derivatives of Exponential and Trigonometric Functions:

We state without proof (at this stage) that:

$$\frac{de^x}{dx} = e^x$$

Actually, this is what make the number e so special. It is the only nonzero function whose derivative is itself.

The derivative of $\sin x$ is:

$$\begin{aligned} \frac{d \sin x}{dx} &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h} \\ &= \sin x \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \lim_{h \rightarrow 0} \frac{\sin h}{h} \\ &= \sin x \lim_{h \rightarrow 0} \frac{1 - 2\sin^2(h/2) - 1}{h} + \cos x \cdot 1 \\ &= \cos x \end{aligned}$$

We can show that

$$\frac{d \cos x}{dx} = -\sin x$$

similarly.

Exercise 6-11: Prove that

$$\begin{aligned} \frac{d \tan x}{dx} &= \sec^2 x \\ \frac{d \cot x}{dx} &= -\csc^2 x \\ \frac{d \sec x}{dx} &= \sec x \tan x \\ \frac{d \csc x}{dx} &= -\csc x \cot x \end{aligned}$$

Exercise 6-12: Evaluate the derivatives of the following functions:

a) $y = \frac{x - \cos x}{x + \cos x}$

b) $y = x^n e^x$

c) $y = \sin 2\theta$

d) $s = \theta^2 \sec \theta$

e) $p = \frac{t^2 \sin t}{t - \tan t}$

Exercise 6-13: Find the following limits:

a) $\lim_{t \rightarrow \pi/4} \frac{\tan t - 1}{t - \pi/4}$

b) $\lim_{t \rightarrow \pi/3} \frac{\cos t - 1/2}{t - \pi/3}$

Chain Rule: If f and g are differentiable then $f(g(x))$ is also differentiable and

$$[f(g(x))]' = f'(g(x)) \cdot g'(x)$$

or more briefly

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Power Rule for Rational Powers: If r is a rational number and $f(x) = x^r$ then $f' = rx^{r-1}$. Let $r = \frac{p}{q}$. Then $y^q = x^p$.

Derivative of both sides gives

$$qy^{q-1}y' = px^{p-1}$$

$$y' = \frac{px^{p-1}}{qx^{\frac{p}{q}(q-1)}} = \frac{p}{q} x^{\frac{p}{q}-1}$$

Exercise 6-14: Find y'

a) $y = (x^2 - 2)^2$

b) $y = [x + (x^3 + 5)^2]^4$

c) $y = \left(t - \frac{1}{t}\right)^2$

d) $y = \sqrt{1 + \sqrt{1 + x}}$

e) $y = te^{-3t^2}$

f) $y = 3^t$

g) $y = 5^{3t}$

h) $y = e^{s^2(1-s)^3}$

i) $y = \cos^2 \theta$

j) $y = 2 \cos^3(5\theta)$

k) $y = \sin(\sin(\pi\theta))$

— End of WEEK —

Review Exercises

Exercise 6-15: Find the derivatives of the following functions:

a) $s = \frac{\sqrt{t}}{3 - 2\sqrt{t}}$

b) $y = x^5 \cos^2(3x^4)$

c) $r = \tan \sqrt{3\theta - 2}$

d) $r = \left(\frac{\cos \theta}{1 - \theta \sin \theta} \right)^2$

e) $p = \frac{1}{(q - q^2 - \cos^2 q)^3}$

f) $y = x^{-5/3} \cos(3x - 1) - \sqrt{2}e^{x-\sqrt{2}}$

Exercise 6-16: Evaluate the derivatives of the following functions:

a) $f = \sin^2 \sqrt{x}$

b) $f = \frac{x}{\cos 4x}$

c) $f = \cos(\sin x)$

d) $f = \sqrt{\cos \sqrt{x}}$

e) $f = x^2 \cos \frac{1}{x}$

f) $f = \frac{1}{\sin^2 x + \cos^2 x}$

Author: Dr. Emre Sermutlu

Last Update: November 9, 2016